

Immunisation Schedule for Children



AGE OF CHILD	VACCINES NEEDED	HOW
At Birth	BCG Bacillus Calmette Guerin	Right arm
	OPV (0) Oral Polio Vaccine	By mouth
6 Weeks	OPV (1) Oral Polio Vaccine	
	RV (1) Rotavirus Vaccine	Intramuscular / Left thigh
	DTaP-IPV // Hib (1) Diphtheria, Tetanus, acellular Pertussis, Inactivated Polio Vaccine and <i>Haemophilus influenzae</i> type b Combined	
	Hep B (1) Hepatitis B Vaccine	
	PCV, (1) Pneumococcal Conjugated Vaccine	Intramuscular / Right thigh
10 Weeks	DTaP-IPV // Hib (2) Diphtheria, Tetanus, acellular Pertussis, Inactivated Polio Vaccine and <i>Haemophilus influenzae</i> type b Combined	Intramuscular / Left thigh
	Hep B (2) Hepatitis B Vaccine	Intramuscular / Right thigh
	14 Weeks	RV (1) Rotavirus Vaccine*
DTaP-IPV // Hib (3) Diphtheria, Tetanus, acellular Pertussis, Inactivated Polio Vaccine and <i>Haemophilus influenzae</i> type b Combined		Intramuscular / Left thigh
Hep B (3) Hepatitis B Vaccine		Intramuscular / Right thigh
PCV, (2) Pneumococcal Conjugated Vaccine		
6 Months	Measles Vaccine (1)	Subcutaneous / Left thigh
9 Months	PCV, (3) Pneumococcal Conjugated Vaccine	Intramuscular / Right thigh
12 Months	Measles Vaccine (2)	Subcutaneous / Right arm
18 Months	DTaP-IPV // Hib (4) Diphtheria, Tetanus, acellular Pertussis, Inactivated Polio Vaccine and <i>Haemophilus influenzae</i> type b Combined	Intramuscular / Left arm
	Td Vaccine Tetanus and reduced strength of diphtheria Vaccine	
6 Years (Boys & girls)	HPV Vaccine	Intramuscular / Left arm
9 Years (Girls only)	HPV Vaccine	Intramuscular / Arm or thigh
12 Years (Boys & girls)	Td Vaccine Tetanus and reduced strength of diphtheria Vaccine	Intramuscular / Left arm

*Rotavirus vaccine should NOT be administered after 24 weeks.

DROPS BY MOUTH
 LIQUID BY MOUTH
 INJECTION

