

Department of Health

IMMUNISATION SCHEDULE FOR CHILDREN

PROFMED



AGE OF CHILD

VACCINES NEEDED

HOW IT'S ADMINISTERED

AT BIRTH

BCG Bacillus Calmette Guerin

● Right arm

OPV (0) Oral Polio Vaccine

● By mouth

6 WEEKS

OPV (1) Oral Polio Vaccine

● By mouth

RV (1) Rotavirus Vaccine

● By mouth

DTaP-IPV // Hib (1) Diphtheria, Tetanus, Acellular Pertussis, Inactivated Polio Vaccine and Haemophilus Influenzae Type b Combined

● Intramuscular / Left thigh

Hep B (1) Hepatitis B Vaccine

● Intramuscular / Right thigh

PCV₇ (1) Pneumococcal Conjugated Vaccine

● Intramuscular / Right thigh

10 WEEKS

DTaP-IPV // Hib (2) Diphtheria, Tetanus, Acellular Pertussis, Inactivated Polio Vaccine and Haemophilus Influenzae Type b Combined

● Intramuscular / Left thigh

Hep B (2) Hepatitis B Vaccine

● Intramuscular / Right thigh

RV (1) Rotavirus Vaccine*

● By mouth

14 WEEKS

DTaP-IPV // Hib (3) Diphtheria, Tetanus, Acellular Pertussis, Inactivated Polio Vaccine and Haemophilus Influenzae Type b Combined

● Intramuscular / Left thigh

Hep B (3) Hepatitis B Vaccine

● Intramuscular / Right thigh

PCV₇ (2) Pneumococcal Conjugated Vaccine

● Intramuscular / Right thigh

6 MONTHS

Measles Vaccine (1)**

● Subcutaneous / Left thigh

9 MONTHS

PCV₇ (3) Pneumococcal Conjugated Vaccine

● Intramuscular / Right thigh

12 MONTHS

Measles Vaccine (2)**

● Subcutaneous / Right arm

18 MONTHS

DTaP-IPV // Hib (4) Diphtheria, Tetanus, Acellular Pertussis, Inactivated Polio Vaccine and Haemophilus Influenzae Type b Combined

● Intramuscular / Left arm

6 YEARS (BOYS & GIRLS)

Td Vaccine Tetanus and Reduced Strength of Diphtheria Vaccine

● Intramuscular / Left arm

9 YEARS (GIRLS ONLY)

HPV Vaccine

● Intramuscular / Arm or thigh

12 YEARS (BOYS & GIRLS)

Td Vaccine Tetanus and Reduced Strength of Diphtheria Vaccine

● Intramuscular / Left arm

*Rotavirus vaccine should NOT be administered after 24 weeks.

**Do not administer with any other vaccine.

● DROPS BY MOUTH

● LIQUID BY MOUTH

● INJECTION